



TEXAS HEART[®] INSTITUTE

The Center for

**WOMEN'S HEART &
VASCULAR HEALTH**

*Supported in part by a generous grant from
Kappa Kappa Gamma Charitable Foundation of Houston*

**Numbers You Need to Know &
Information to Share With Your Doctor**

Numbers You Need to Know

“My goal is first, to alert women to the fact that they definitely are at risk for heart disease, and...it is critical that women keep an eye on their numbers!” —Dr. Stephanie Coulter

This worksheet provides a place for you to record your numbers. See the following page for a brief explanation of these common blood tests and measurements performed by your doctor. Refer to the back page for information to share with your doctor.

Blood Pressure

_____ / _____
systolic / diastolic

	systolic	AND	diastolic
Normal:	<120	AND	<80
Elevated:	120–129	AND	<80
High Blood Pressure (Hypertension)			
Stage 1:	130–139	OR	80–89
Stage 2:	≥140	OR	≥90
Hypertensive Crisis	>180	AND/OR	>120

Waist Circumference (in.)

High-Risk Levels

Men:	>40 inches
Women:	>35 inches

Fasting Glucose

Normal:	60–99
Pre-diabetes:	100–125
Diabetes:	>126

Body Mass Index (BMI)

Height (in) _____
Weight (lbs) _____

BMI Equation: $\frac{\text{Weight in pounds} \times 703}{\text{Height in inches} \times \text{Height in inches}}$

Underweight:	<18.5
Normal:	18.5–24.9
Overweight:	25.0–29.9
Obese:	>30.0

Total Cholesterol

Desirable:	<200
Borderline high:	200–239
High:	≥240

HDL (“Good” Cholesterol)

Low:	<40
Normal:	40–59
Optimal:	≥60

LDL (“Bad” Cholesterol)

Optimal:	<100
Near optimal:	100–129
Borderline high:	130–159
High:	≥160
Very high:	>190

Total Cholesterol/HDL Ratio

Desirable:	<4.6
-------------------	----------------

Fasting Triglycerides

Normal:	<150
Borderline high:	150–199
High:	≥200

Definitions of Numbers You Need to Know

This worksheet briefly explains the common blood tests and measurements taken by your doctor. Use the space on the previous page to record your numbers.

Blood Pressure. When your heart beats, it pumps blood through blood vessels called arteries and creates pressure in them. The higher number (systolic) represents the pressure while your heart is beating. The lower number (diastolic) represents the pressure when your heart is resting between beats. Your blood pressure can change from minute to minute when you change your position or during physical activities, or when you are stressed or sleeping. High blood pressure can damage the walls of your arteries. If your first reading is high, your doctor may take several more readings before deciding whether your blood pressure may be a health concern.

Waist Circumference. A waist circumference measures belly fat. Too much belly fat is associated with a greater health risk than fat located in the hips, thighs, or buttocks. Too much belly fat is an “independent risk factor” for high cholesterol and heart disease, regardless of your Body Mass Index or BMI score, which could be normal. A large waist circumference is also associated with an increased risk for type 2 diabetes and hypertension.

Fasting Glucose. A fasting glucose test shows the amount of sugar in your blood. If you have eaten within 8 hours before this test, your glucose level may be falsely elevated. Glucose rises rapidly after meals but should return to near the fasting value within 2 hours. A high blood glucose level may be a sign of diabetes or pre-diabetes.

Body Mass Index. Body Mass Index (BMI), which is a number calculated from your height and weight, is used to estimate body fat. The accuracy of the BMI in determining your body fat is influenced by factors such as your fitness level, muscle mass, bone structure, sex, and ethnic origin. This means that some individuals may have a high BMI but not have a high percentage of body fat. However, BMI provides a reliable indicator of body fat for most people. Therefore, it is used to screen for weight problems that may lead to health concerns.

Total Cholesterol. Cholesterol is a measure of the amount of fat in your blood stream. There is good fat (HDL, high-density lipoprotein) and bad fat (LDL, low-density lipoprotein). HDL number should be “high”; LDL number should be “low”. Your body produces all the cholesterol it needs; however, cholesterol is also added to your body from the foods you eat—mainly saturated fats. Foods rich in saturated fat include butter fat in milk products, fat from red meat, and tropical oils such as coconut oil. If you keep your total cholesterol level within the desirable range, you may reduce your chances of heart attack and stroke.

HDL Cholesterol. HDL cholesterol, or “good” cholesterol, helps clear away bad fatty substances that can stick to your arteries and damage them. The higher the concentration of HDL in your blood, the less likely you are to develop coronary heart disease. The amount of HDL cholesterol you have may be increased in a number of ways, including getting regular physical activity, not smoking, and losing weight.

LDL Cholesterol. LDL cholesterol, or “bad” cholesterol, is a major contributor to blockages in the arteries that can cause heart attack, stroke, and poor circulation. The higher your LDL concentration, the more likely you are to develop these conditions. LDL cholesterol may be decreased by reducing trans fats and saturated fats that you eat and by eating more fiber-rich foods. If you have eaten within 12 hours before having this test, your LDL level may be incorrect.

Total Cholesterol/HDL Ratio. The total cholesterol/HDL ratio is an indicator of your potential for developing blockages in the arteries of your heart. A ratio greater than 4.5 is considered a high risk for coronary heart disease. The ratio may be decreased by increasing your good (HDL) cholesterol and/or decreasing your bad (LDL) cholesterol.

Fasting Triglycerides. Triglycerides are in a class of blood fats known as “lipids.” If you have eaten within 12 hours of having your blood tested for triglycerides, your level may be falsely high. A high triglyceride level can result in hardening of the arteries, a condition often called “coronary heart disease.” High levels of triglycerides are more common in people who are overweight, have diabetes, and/or who do not process fat properly. Your triglyceride levels may be reduced by decreasing your intake of simple sugars (sweets) and alcohol and by increasing your physical activity.

Information to Share With Your Doctor

This worksheet of questions helps you prepare for a doctor's visit. Share your answers with your doctor as a reminder to you both of other important information that is helpful in assessing your risk.

- At what age did you begin menopause, if you have? _____
- Did you have a hysterectomy? Yes No
- Were the ovaries removed? Yes No
 - If so, when? _____
- Did you take hormones or hormone replacement therapy? Yes No
 - For how many years? _____
 - Did you take estrogen alone (for example, Premarin)? Yes No
 - Did you take estrogen and progesterone (for example, Prempro)? Yes No
- Do you have diabetes? Yes No
- Do you take insulin? Yes No
- Do you have a history of gestational diabetes? Yes No
- Do you have a history of high blood pressure during pregnancy (Preeclampsia)? Yes No
- Do you have high blood pressure? Yes No
 - Do you take medicine for high blood pressure? Yes No
- Do you take medicine for high cholesterol? Yes No
- Do you smoke? Yes No
 - Do you use tobacco daily? Yes No
 - If you no longer smoke, how long has it been since you quit? _____
- Do you have any family history of coronary artery disease or peripheral vascular disease? Yes No
 - If so, which male relatives under the age of 55? _____
 - If so, which female relatives under the age of 65? _____
- During the past two weeks:
 - Have you often been bothered by feeling down, depressed or hopeless? Yes No
 - Have you often been bothered by little interest or pleasure in doing things? Yes No
 - Have you ever been diagnosed with Lupus or Rheumatoid Arthritis? Yes No

Additional Notes:
